



THE JOURNAL OF  
**CLINICAL PSYCHIATRY**  
THE OFFICIAL JOURNAL OF THE AMERICAN SOCIETY OF CLINICAL PSYCHOPHARMACOLOGY

## **Supplementary Material**

**Article Title:** A Randomized, Double-Blind, Placebo-Controlled Trial of Aripiprazole Lauroxil in Acute Exacerbation of Schizophrenia

**Authors:** Herbert Y. Meltzer, MD; Robert Risinger, MD; Henry A. Nasrallah, MD; Yangchun Du, PhD; Jacqueline Zummo, MPH, MBA; Lisa Corey, MS; Anjana Bose, PhD; Srdjan Stankovic, MD, MSPH; Bernard L. Silverman, MD; and Elliot W. Ehrich, MD

**DOI Number:** 10.4088/JCP.14m09741

### **List of Supplementary Material for the article**

1. [eTable 1](#) Change from baseline in PANSS total score using Full Analysis Set (FAS)
2. [eFigure 1](#) Incidence of injection site pain

### **Disclaimer**

This Supplementary Material has been provided by the author(s) as an enhancement to the published article. It has been approved by peer review; however, it has undergone neither editing nor formatting by in-house editorial staff. The material is presented in the manner supplied by the author.

© Copyright 2015 Physicians Postgraduate Press, Inc.

Supplementary eTable 1. Change from baseline in PANSS total score using Full Analysis Set (FAS)

	Aripiprazole lauroxil		
Parameter	441 mg (N=196)	882 mg (N=204)	Placebo (N=196)
LOCF Analysis			
N	196	204	196
Baseline, mean (SD)	92.6 (10.2)	92.0 (10.8)	93.9 (11.3)
Change from Baseline at Day 85: LS Mean (SE)	-20.9 (1.4)	-21.8 (1.4)	-9.8 (1.4)
Adjusted LS mean (SE) difference from placebo <sup>a</sup>	-10.9 (1.8)	-11.9 (1.8)	
Adjusted p-value vs. placebo	<0.001	<0.001	
MMRM			
N	196	204	196
Change from Baseline at Day 85: LS mean (SE)	-22.3 (1.5)	-22.9 (1.4)	-10.6 (1.6)
LS mean (SE) difference from placebo	-11.8 (2.1)	-12.4 (2.1)	
p-value vs. placebo	<0.001	<0.001	
MMRM for subset of severe population			

N	95	100	99
Baseline, mean (SD)	101.3 (6.0)	101.0 (6.4)	102.7 (7.9)
Change from Baseline at Day 85	-22.14 (2.4)	-24.05 (2.3)	-7.44 (2.8)
LS mean (SE)			
LS mean (SE) difference from placebo	-14.7 (3.5)	-16.6 (3.4)	
p-value vs. placebo	<0.001	<0.001	

<sup>a</sup>Adjusted LS Mean (SE) difference controlling for overall Type I error rate<sup>b</sup>

<sup>b</sup>Cui L, Hung HMJ, Wang S. Modification of sample size in group sequential clinical trials. *Biometrics*. 1999;55 (3):853-857.

### Supplementary eFigure 1. Incidence of injection site pain

